

Survey Question: Did you use any of these drugs when you were pregnant? Marijuana (pot, bud) or hashish (hash), Amphetamines (uppers, ice, speed, crystal meth, crank), Cocaine (rock, coke, crack) or heroin (smack, horse), Tranquilizers (downers, ludes) or hallucinogens (LSD/acid, PCP/angel dust, ecstasy), Sniffing gasoline, glue, hairspray, or other aerosols

Year	State of Hawai'i				Native Hawaiian			
	No.	Illicit drug use during pregnancy	95%CI Lower Limit	95%CI Upper Limit	No.	Illicit drug use during pregnancy	95%CI Lower Limit	95%CI Upper Limit
2000	500	2.8%	2.1%	3.6%	200	4.5%	3.4%	5.9%
2001	600	3.4%	2.6%	4.3%	200	3.8%	2.9%	5.1%
2002	300	2.0%	1.5%	2.7%	200	4.4%	3.2%	6.0%
2003	500	2.9%	2.2%	3.9%	200	5.1%	3.8%	6.8%
2004	500	3.0%	2.3%	4.0%	200	4.2%	2.6%	6.7%
2005	400	2.3%	1.7%	3.3%	**	**	**	**
2006	500	2.9%	2.1%	3.9%	200	4.7%	2.9%	7.6%
2007	500	2.8%	2.1%	3.8%	200	4.2%	2.5%	6.8%
2008	400	2.3%	1.7%	3.3%	200	3.7%	2.2%	6.2%
2009	500	3.0%	2.1%	4.2%	200	4.2%	2.5%	7.1%
2010	400	2.5%	1.7%	3.7%	**	**	**	**
2011	600	3.4%	2.5%	4.6%	300	5.3%	3.3%	8.3%

**The estimate has been suppressed because 1) The relative standard error is greater than 50% or when the relative standard error can't be determined. Consider aggregating years to decrease the relative standard error and improve the reliability of the estimate. 2) the observed number of events is very small and not appropriate for publication, or 3) it could be used to calculate the number in a cell that has been suppressed.

Source: Hawai'i State, Department of Health, Hawai'i Health Data Warehouse. Hawai'i Pregnancy Risk Assessment Monitoring System, PRAMS Health Indicator Report- Illicit drug use in pregnancy