

Kūpuna 5B5

Native Hawaiian Adults Aged 50-75 have had a Colorectal Screening n Hawai'i: 2011-2014

Survey Question: Persons aged 50 to 75 years who have had a blood stool test in the past year, sigmoidoscopy in the past 5 years and blood stool test in the past 3 years, or a colonoscopy in the past 10 years?

Year	State of Hawai'i						
	Yes			No			Estimated Population
	#	%	CI (95%)	#	%	CI (95%)	#
2011	212,100	59.4	57.1-61.7	144,900	40.6	38.3-42.9	357,000
2012	246,100	61.1	58.6-63.7	156,500	38.9	36.3-41.4	402,600
2013	256,200	66.4	64.1-68.7	129,700	33.6	31.3-35.9	385,800
2014	257,200	66.5	64.2-68.8	129,700	33.5	31.2-35.8	386,900

Year	Native Hawaiian						
	Yes			No			Estimated Population
	#	%	CI (95%)	#	%	CI (95%)	#
2011	16,100	53.4	46.5-60.2	14,000	46.6	39.8-53.5	30,100
2012	17,900	54.7	47.3-62.2	14,800	45.3	37.8-52.7	32,700
2013	21,400	62.8	56.0-69.5	12,700	37.2	30.5-44.0	34,000
2014	22,600	60.4	54.4-66.4	14,800	39.6	33.6-45.6	37,400

Source: Hawai'i State, Department of Health, Office of Health Status Monitoring, Hawai'i Health Data Warehouse, Hawai'i Behavioral Risk Factor Surveillance System, Colorectal screening (50-75), for the State of Hawai'i, for the Year(s)-2011, 2012, 2013, 2014