

‘Ōpio 3B28

Native Hawaiian Public High School Students Who Used Birth Control Pills, Condoms, Depo-Provera, Nuva Ring, Implanon, or any IUD, to Prevent Pregnancy, Among Students Who had Sexual Intercourse During the Past 3 Months in Hawai‘i: 2000-2015

Survey Question: The last time you had sexual intercourse, what one method did you or your partner use to prevent pregnancy?

| Year | Sex (current)-used birth control last intercourse | | | | | |
|------|---|-------|-----------|-----------------|-------|-----------|
| | State of Hawai‘i | | | Native Hawaiian | | |
| | # | % | CI | # | % | CI |
| 2005 | 6,300 | 56.9% | 48.0-65.9 | n/r | n/r | n/r |
| 2007 | 6,400 | 60.1% | 51.9-68.4 | n/r | n/r | n/r |
| 2009 | 7,000 | 56.4% | 48.7-64.1 | n/r | n/r | n/r |
| 2011 | 4,700 | 52.7% | 49.3-56.1 | 1,400 | 55.6% | 49.2-62.0 |
| 2013 | 5,200 | 60.5% | 55.6-65.3 | 1,500 | 58.8% | 48.7-68.9 |
| 2015 | 5,000 | 64.5% | 58.0-70.9 | 1,300 | 55.1% | 45.9-64.3 |

Source: Hawai‘i State, Department of Health, Hawai‘i Health Data Warehouse, Youth Risk Behavioral Survey, Current Sexual Activity in Hawai‘i