

Table 3.23 Water Quality at Public Beaches in Hawai'i by Island: 2004-2008.

Year	Enterococci Density ^a	Island								
		Hawai'i			Maui	Lāna'i	Moloka'i	O'ahu	Kauai	Total
		Hawai'i	Hilo Shoreline	Kona Shoreline						
2004	Lowest	0.6	0.6	1.2	1.2	(N)	(N)	0.3	1.0	0.3
	Mean	3.7	4.7	2.7	2.9	(N)	(N)	4.5	5.4	4.1
	Highest	230.0	230.0	7.3	18.2	(N)	(N)	110.7	227.0	230.0
2005	Lowest	0.3	0.9	0.3	0.5	(N)	(N)	0.4	0.3	0.3
	Mean	3.9	5.2	2.2	2.1	(N)	(N)	5.1	5/ 4.6	4.2
	Highest	15.7	15.7	8.1	27.0	(N)	(N)	47.3	243.9	243.9
2006	Lowest	1.0	0.5	1.0	1.8	(N)	(N)	0.7	0.6	0.5
	Mean	2.8	6.0	2.0	3.5	(N)	(N)	4.6	6.5	4.6
	Highest	20.1	18.1	20.0	6.1	(N)	(N)	80.0	144.9	144.9
2007	Lowest	2.3	2.3	2.3	2.3	(N)	(N)	2.3	2.3	2.3
	Mean	3.7	4.1	3.2	3.4	(N)	(N)	3.9	4.6	3.8
	Highest	4.5	75.0	4.5	22.6	(N)	(N)	475.7	700.0	700.0
2008	Lowest	2.3	2.5	2.3	2.7	(N)	(N)	2.3	2.3	2.3
	Mean	4.6	4.8	4.1	3.6	(N)	(N)	4.7	5.0	4.5
	Highest	64.0	10.6	64.0	4.2	(N)	(N)	137.0	99.9	137.0

^a Geometric mean, number per 100 ml. The geometric mean standard for Enterococci density is 7 per 100 ml.
(N) Not applicable.

Source: Hawai'i State, Department of Health, Clean Water Branch, records.