Table 7.148

Men Who Had a Prostate-Specific Antigen (PSA) Test (Among Men Age 40+) by Race-Ethnicity in Hawai'i: 2013

| Race-Ethnicity | All Men Aged 40+ | | | |
|------------------------------------|--|------------|---|------------|
| | Ever Had a Prostate-Specific Antigen (PSA) Test | | Prostate-Specific Antigen (PSA) Test within Past 2 Years | |
| | % | C.I. (95%) | % | C.I. (95%) |
| Caucasian | 49.3 | 44.7-53.9 | 34.4 | 30.0-38.8 |
| Native Hawaiian | 27.0 | 19.3-34.6 | 19.1 | 12.3-26.0 |
| Chinese | 44.4 | 31.4-57.4 | 27.7 | 16.3-39.1 |
| Filipino | 26.1 | 18.9-33.4 | 20.2 | 13.6-26.8 |
| Japanese | 45.8 | 39.8-51.8 | 34.1 | 28.5-39.6 |
| Black | n/r | n/r | n/r | n/r |
| Native Alaskan/ American Indian | n/r | n/r | n/r | n/r |
| Other Asian | n/r | n/r | n/r | n/r |
| Other Pacific Islander | n/r | n/r | n/r | n/r |
| Other | 40.2 | 16.6-63.7 | 35.8 | 11.5-60.1 |
| Total | 40.8 | 37.9-43.7 | 29.5 | 26.8-32.2 |

Source: Hawai'i State, Department of Health, Office of Health Status Monitoring. Hawaii Health Data Warehouse. Behavioral Risk Factor Surveillance System.