

Table 7.149

Native Hawaiian Men Who Ever Had a Prostate-Specific Antigen (PSA) Test (Among Men Age 40+) in Hawai'i: 2001-2013

Year	Native Hawaiian Men Aged 40+			
	Ever Had a Prostate-Specific Antigen (PSA) Test		Prostate-Specific Antigen (PSA) Test within Past 2 Years	
	%	C.I. (95%)	%	C.I. (95%)
2013	27.0	19.3-34.6	19.1	12.3-26.0
2012	27.9	18.6-37.2	15.7	9.4-21.9
2011	37.1	28.1-46.1	27.3	19.2-35.5
2010	51.8	42.1-61.4	38.6	29.6-47.7
2009	38.3	25.3-51.3	32.5	19.7-45.3
2008	44.4	34.6-54.1	38.0	28.5-47.6
2007	46.5	37.4-55.7	32.2	24.2-40.2
2006	31.0	23.0-39.0	27.4	19.6-35.1
2005	38.8	24.0-53.6	35.6	21.0-50.3
2004	35.4	19.6-51.1	30.6	15.2-46.0
2003	42.0	28.0-56.0	39.5	25.6-53.3
2002	39.7	28.6-50.8	33.6	22.8-44.5
2001	48.0	35.4-60.6	36.9	24.6-49.2

**Source:** Hawai'i State, Department of Health, Office of Health Status Monitoring. Hawaii Health Data Warehouse. Behavioral Risk Factor Surveillance System.