

Table 7.151

Native Hawaiian Adults Who Had a Blood Stool Test Using a Home Kit (Among Adults Age 50+) in Hawai'i: 2001-2013

Year	Native Hawaiian Men Aged 50+			
	Blood Stool Test Done Using a Home Kit		Most Recent Blood Stool Test Done Using a Home Kit within 1 Year	
	%	C.I. (95%)	%	C.I. (95%)
2013	44.7	37.7-51.6	24.1	17.3-30.9
2012	35.4	28.9-41.9	13.6	9.0-18.3
2011	48.0	41.6-54.5	18.0	12.9-23.2
2010	42.8	36.5-49.1	18.7	13.5-23.8
2009	45.3	35.6-55.0	15.0	7.6-22.4
2008	40.7	33.9-47.5	15.8	11.1-20.6
2007	47.4	40.9-53.9	20.7	15.4-25.9
2006	42.6	35.6-49.6	20.0	14.1-26.0
2005	42.7	32.4-53.0	23.1	14.5-31.6
2004	41.4	27.8-55.0	16.1	3.7-28.4
2003	52.7	42.4-63.0	31.8	22.4-41.2
2002	39.9	31.2-48.5	20.4	14.0-26.8
2001	36.2	26.6-45.9	24.4	15.4-33.5

Source: Hawai'i State, Department of Health, Office of Health Status Monitoring. Hawaii Health Data Warehouse. Behavioral Risk Factor Surveillance System.