

Table 7.152 Native Hawaiians Blood Stool Test Using a Home Kit (50+) in Hawai'i: 2001-2013

Survey Question: A blood stool test is a test that may use a special kit at home to determine whether the stool contains blood. Have you ever had this test using a home kit?

| Year | Native Hawaiian | | | | | | Estimated Population No. |
|------|-----------------|------|------------|--------|------|------------|-----------------------------|
| | Yes | | | No | | | |
| | No. | % | C.I. (95%) | No. | % | C.I. (95%) | |
| 2013 | 16,800 | 44.7 | 37.7-51.6 | 20,800 | 55.3 | 48.4-62.3 | 37,500 |
| 2012 | 14,400 | 35.4 | 28.9-41.9 | 26,200 | 64.6 | 58.1-71.1 | 40,600 |
| 2011 | 16,900 | 48.0 | 41.6-54.5 | 18,300 | 52.0 | 45.5-58.4 | 35,200 |
| 2010 | 18,400 | 42.8 | 36.5-49.1 | 24,600 | 57.2 | 50.9-63.5 | 43,100 |
| 2009 | 16,400 | 45.3 | 35.6-55.0 | 19,800 | 54.7 | 45.0-64.4 | 36,200 |
| 2008 | 17,100 | 40.7 | 33.9-47.5 | 25,000 | 59.3 | 52.5-66.1 | 42,200 |
| 2007 | 21,200 | 47.4 | 40.9-53.9 | 23,500 | 52.6 | 46.1-59.1 | 44,700 |
| 2006 | 17,600 | 42.6 | 35.6-49.6 | 23,700 | 57.4 | 50.4-64.4 | 41,300 |
| 2005 | 19,100 | 42.7 | 32.4-53.0 | 25,600 | 57.3 | 47.0-67.6 | 44,800 |
| 2004 | 17,300 | 41.4 | 27.8-55.0 | 24,400 | 58.6 | 45.0-72.2 | 41,700 |
| 2003 | 15,600 | 52.7 | 42.4-63.0 | 13,900 | 47.3 | 37.0-57.6 | 29,500 |
| 2002 | 14,100 | 39.9 | 31.2-48.5 | 21,200 | 60.1 | 51.5-68.8 | 35,300 |
| 2001 | 13,600 | 36.2 | 26.6-45.9 | 23,900 | 63.8 | 54.1-73.4 | 37,500 |

Source: Hawai'i State, Department of Health, Office of Health Status Monitoring. Hawaii Health Data Warehouse. Behavioral Risk Factor Surveillance System. Blood stool test-using a home kit (50+), for the State of Hawai'i

| Year | State of Hawai'i | | | | | | |
|------|------------------|------|------------|---------|------|------------|----------------------|
| | Yes | | | No | | | Estimated Population |
| | No. | % | C.I. (95%) | No. | % | C.I. (95%) | No. |
| 2013 | 201,400 | 43.2 | 41.0-45.4 | 264,900 | 56.8 | 54.6-59.0 | 466,400 |
| 2012 | 215,400 | 44.6 | 42.3-46.8 | 268,100 | 55.4 | 53.2-57.7 | 483,600 |
| 2011 | 202,200 | 45.6 | 43.6-47.7 | 240,800 | 54.4 | 52.3-56.4 | 443,000 |
| 2010 | 205,100 | 47.2 | 45.3-49.2 | 229,400 | 52.8 | 50.8-54.7 | 434,500 |
| 2009 | 203,600 | 49.0 | 46.2-51.8 | 212,000 | 51.0 | 48.2-53.8 | 415,500 |
| 2008 | 211,500 | 49.7 | 47.6-51.8 | 214,300 | 50.3 | 48.2-52.4 | 425,800 |
| 2007 | 207,000 | 49.5 | 47.3-51.7 | 211,100 | 50.5 | 48.3-52.7 | 418,100 |
| 2006 | 190,800 | 46.9 | 44.7-49.1 | 216,200 | 53.1 | 50.9-55.3 | 407,000 |
| 2005 | 188,100 | 47.5 | 44.3-50.8 | 207,700 | 52.5 | 49.2-55.7 | 395,800 |
| 2004 | 167,300 | 45.0 | 40.9-49.2 | 204,300 | 55.0 | 50.8-59.1 | 371,600 |
| 2003 | 205,200 | 54.7 | 51.9-57.6 | 169,600 | 45.3 | 42.4-48.1 | 374,800 |
| 2002 | 150,300 | 43.7 | 41.2-46.3 | 193,400 | 56.3 | 53.7-58.8 | 343,700 |
| 2001 | 170,100 | 49.6 | 46.4-52.9 | 172,600 | 50.4 | 47.1-53.6 | 342,700 |

Source: Hawai'i State, Department of Health, Office of Health Status Monitoring. Hawaii Health Data Warehouse. Behavioral Risk Factor Surveillance System. Blood stool test-using a home kit (50+), for the State of Hawai'i