

Table 7.153

Native Hawaiians who have had a blood stool test in the past year, sigmoidoscopy in the past 5 years and blood stool test in the past 3 years, or a colonoscopy in the past 10 years in Hawai'i: 2011-2013

Year	Native Hawaiian						
	Yes			No			Estimated Population
	No.	%	C.I.(95%)	No.	%	C.I.(95%)	No.
2013	12,700	37.2	30.5-44.0	21,400	62.8	56.0-69.5	34,000
2012	15,600	44.2	37.1-51.3	19,800	55.8	48.7-62.9	35,400
2011	14,000	46.6	39.8-53.5	16,100	53.4	46.5-60.2	30,100

Year	State of Hawai'i						
	Yes			No			Estimated Population
	No.	%	C.I.(95%)	No.	%	C.I.(95%)	No.
2013	129,700	33.6	31.3-35.9	256,200	66.4	64.1-68.7	385,800
2012	156,500	38.9	36.3-41.4	246,100	61.1	58.6-63.7	402,600
2011	144,900	40.6	38.3-42.9	212,100	59.4	57.1-61.7	357,000

Source: Hawai'i State, Department of Health, Office of Health Status Monitoring. Hawaii Health Data Warehouse. Behavioral Risk Factor Surveillance System. Colorectal screening (50-75), for the State of Hawai'i