

**Table 7.154** Sigmoidoscopy or a Colonoscopy (Among Adults Age 50+) by Race-Ethnicity in Hawai'i: 2013

Race-Ethnicity	All Men Aged 50+			
	Had a Sigmoidoscopy or a Colonoscopy		Most Recent Sigmoidoscopy or Colonoscopy within 1 Year	
	%	C.I. (95%)	%	C.I. (95%)
Caucasian	70.2	67.2-73.2	16.6	13.9-19.4
Native Hawaiian	58.7	51.8-65.5	16.2	10.7-21.7
Chinese	68.2	59.2-77.2	8.9	4.2-13.5
Filipino	53.0	45.7-60.3	14.6	9.3-20.0
Japanese	77.9	74.4-81.4	17.7	14.0-21.3
Black	n/r	n/r	n/r	n/r
Native Alaskan/ American Indian	n/r	n/r	n/r	n/r
Other Asian	76.8	64.9-88.7	19.7	4.4-35.0
Other Pacific Islander	n/r	n/r	n/r	n/r
Other	41.0	24.4-57.5	12.3	1.1-23.4
<b>Total</b>	68.3	66.2-70.4	15.9	14.1-17.6

**Source:** Hawai'i State, Department of Health, Office of Health Status Monitoring. Hawaii Health Data Warehouse. Behavioral Risk Factor Surveillance System.