

Table 7.155

Native Hawaiian Adults Who Had a Sigmoidoscopy or a Colonoscopy (Among Adults Age 50+) in Hawai'i: 2001-2013

| Year | Native Hawaiian Men Aged 50+ | | | |
|------|--------------------------------------|------------|--|------------|
| | Had a Sigmoidoscopy or a Colonoscopy | | Most Recent Sigmoidoscopy or Colonoscopy within 1 Year | |
| | % | C.I. (95%) | % | C.I. (95%) |
| 2013 | 58.7 | 51.8-65.5 | 16.2 | 10.7-21.7 |
| 2012 | 56.1 | 49.2-63.0 | 15.9 | 10.0-21.7 |
| 2011 | 51.7 | 45.2-58.1 | 15.7 | 11.0-20.4 |
| 2010 | 54.0 | 47.4-60.6 | 47.4 | 40.7-54.1 |
| 2009 | 54.7 | 45.1-64.4 | 47.9 | 38.1-57.6 |
| 2008 | 51.5 | 44.3-58.6 | 49.1 | 41.9-56.4 |
| 2007 | 48.8 | 42.2-55.3 | 51.2 | 44.7-57.8 |
| 2006 | 42.9 | 35.8-50.1 | 57.1 | 49.9-64.2 |
| 2005 | 46.3 | 35.8-56.8 | 53.7 | 43.2-64.2 |
| 2004 | 27.0 | 15.8-38.1 | 73.0 | 61.9-84.2 |
| 2003 | 39.2 | 28.9-49.4 | 60.8 | 50.6-71.1 |
| 2002 | 34.0 | 25.7-42.3 | 66.0 | 57.7-74.3 |
| 2001 | 34.9 | 25.7-44.2 | 65.1 | 55.8-74.3 |

Source: Hawai'i State, Department of Health, Office of Health Status Monitoring. Hawaii Health Data Warehouse. Behavioral Risk Factor Surveillance System.