Table 7.178

**Survey Question:** During the past 12 months, have you had either a seasonal flu shot or a seasonal flu vaccine that was sprayed in your nose?

Year	Native Hawaiian Adults 65+									
	Yes			No			Estimated Population			
	No.	%	C.I. (95%)	No.	%	C.I. (95%)	No.			
2013	8,400	68.8	59.6-78.1	3,800	31.2	21.9-40.4	12,200			
2012	9,200	61.5	50.1-73.0	5,800	38.5	27.0-49.9	15,000			
2011	10,000	63.6	54.0-73.3	5,700	36.4	26.7-46.0	15,700			

Year	State of Hawai'i Adults 65+									
	Yes			No			Estimated Population			
	No.	%	C.I. (95%)	No.	%	C.I. (95%)	No.			
2013	145,000	69.9	67.1-72.8	62,300	30.1	27.2-32.9	207,300			
2012	130,900	62.7	59.5-65.9	77,800	37.3	34.1-40.5	208,700			
2011	125,600	64.7	61.9-67.5	68,600	35.3	32.5-38.1	194,200			

**Source:** Hawai'i State, Department of Health, Office of Health Status Monitoring. Hawaii Health Data Warehouse. Behavioral Risk Factor Surveillance System. Flu-had shot or spray in past 12 months (65+), for the State of Hawai'i.