

Table 7.181 updated

Native Hawaiian Bad Mental Health During the Past 30 Days in Hawai'i: 2000-2014

| Year | Native Hawaiian | | | | | | | | | | | | Estimated Population |
|------|-----------------|------|------------|-------------|------|------------|--------------|------|------------|-----------------|------|------------|----------------------|
| | Zero days | | | 1 to 6 days | | | 7 to 13 days | | | 14 or more days | | | |
| | No. | % | C.I. (95%) | No. | % | C.I. (95%) | No. | % | C.I. (95%) | No. | % | C.I. (95%) | |
| 2014 | 81,400 | 61.5 | 57.0-66.0 | 24,700 | 18.7 | 15.0-22.4 | 13,900 | 10.5 | 8.3-12.8 | 12,300 | 9.3 | 6.0-12.5 | 132,200 |
| 2013 | 70,300 | 62.4 | 58.1-66.7 | 18,500 | 16.4 | 13.3-19.5 | 8,600 | 7.6 | 4.8-10.4 | 15,400 | 13.6 | 10.6-16.7 | 112,800 |
| 2012 | 83,500 | 66.5 | 61.7-71.3 | 20,400 | 16.2 | 12.6-19.8 | 12,800 | 10.2 | 7.7-12.7 | 9,000 | 7.2 | 3.9-10.4 | 125,700 |
| 2011 | 71,700 | 65.2 | 60.4-70.0 | 17,100 | 15.5 | 11.7-19.4 | 6,300 | 5.7 | 3.5-7.9 | 14,900 | 13.5 | 10.1-16.9 | 110,000 |
| 2010 | 90,400 | 69.6 | 64.6-74.6 | 21,400 | 16.5 | 12.4-20.6 | 4,900 | 3.8 | 1.9-5.7 | 13,100 | 10.1 | 6.8-13.4 | 129,800 |
| 2009 | 79,600 | 64.9 | 59.8-70.0 | 20,600 | 16.8 | 13.0-20.6 | 10,200 | 8.4 | 4.6-12.1 | 12,200 | 9.9 | 7.1-12.7 | 122,600 |
| 2008 | 90,700 | 65.7 | 61.0-70.4 | 18,700 | 13.6 | 10.5-16.7 | 9,700 | 7.1 | 4.4-9.8 | 18,800 | 13.7 | 10.5-16.8 | 138,000 |
| 2007 | 92,800 | 68.6 | 64.4-72.8 | 24,300 | 18.0 | 14.4-21.5 | 6,800 | 5.1 | 3.2-7.0 | 11,400 | 8.4 | 6.1-10.7 | 135,400 |
| 2006 | 80,000 | 67.3 | 62.6-72.1 | 16,300 | 13.7 | 10.5-16.9 | 8,600 | 7.3 | 4.1-10.4 | 13,900 | 11.7 | 8.5-14.9 | 118,800 |
| 2005 | 70,400 | 60.6 | 55.7-65.5 | 25,000 | 21.5 | 17.4-25.6 | 5,500 | 4.8 | 2.9-6.6 | 15,300 | 13.1 | 9.5-16.8 | 116,200 |
| 2004 | 75,800 | 62.5 | 53.9-71.0 | 20,800 | 17.1 | 10.3-23.9 | 8,000 | 6.6 | 1.9-11.3 | 16,900 | 13.9 | 7.7-20.1 | 121,400 |
| 2003 | 62,900 | 77.5 | 72.3-82.6 | 9,600 | 11.8 | 7.8-15.8 | 1,900 | 2.4 | 1.0-3.8 | 6,800 | 8.4 | 4.8-11.9 | 81,200 |
| 2002 | 85,100 | 75.3 | 71.1-79.5 | 12,800 | 11.4 | 7.9-14.8 | 5,500 | 4.9 | 3.2-6.6 | 9,600 | 8.5 | 6.1-10.9 | 113,000 |
| 2001 | 98,100 | 85.6 | 81.9-89.3 | 8,800 | 7.7 | 4.8-10.5 | 3,100 | 2.7 | 0.9-4.5 | 4,600 | 4.0 | 2.2-5.8 | 114,600 |
| 2000 | 97,000 | 69.8 | 65.8-73.9 | 21,100 | 15.2 | 12.1-18.3 | 10,100 | 7.2 | 5.0-9.5 | 10,700 | 7.7 | 5.4-10.0 | 138,900 |

Source: Hawai'i State, Department of Health, Office of Health Status Monitoring. Hawaii Health Data Warehouse. Behavioral Risk Factor Surveillance System. Health-mental bad past 30 days

| Year | State of Hawai'i | | | | | | | | | | | | Estimated Population |
|------|------------------|------|------------|-------------|------|------------|--------------|-----|------------|-----------------|-----|------------|----------------------|
| | Zero days | | | 1 to 6 days | | | 7 to 13 days | | | 14 or more days | | | |
| | No. | % | C.I. (95%) | No. | % | C.I. (95%) | No. | % | C.I. (95%) | No. | % | C.I. (95%) | |
| 2014 | 780,500 | 70.6 | 69.1-72.2 | 184,400 | 16.7 | 15.4-18.0 | 93,100 | 8.4 | 7.5-9.3 | 47,000 | 4.3 | 3.6-4.9 | 1,105,100 |
| 2013 | 789,100 | 72.5 | 71.0-73.9 | 160,800 | 14.8 | 13.6-16.0 | 49,600 | 4.6 | 3.9-5.3 | 89,400 | 8.2 | 7.4-9.0 | 1,088,900 |
| 2012 | 749,700 | 70.2 | 68.6-71.8 | 176,900 | 16.6 | 15.2-17.9 | 50,500 | 4.7 | 3.9-5.5 | 90,700 | 8.5 | 7.6-9.4 | 1,067,800 |
| 2011 | 735,900 | 70.6 | 68.9-72.2 | 170,200 | 16.3 | 14.9-17.7 | 48,200 | 4.6 | 3.9-5.3 | 88,800 | 8.5 | 7.5-9.5 | 1,043,000 |
| 2010 | 718,600 | 72.0 | 70.3-73.7 | 166,200 | 16.6 | 15.2-18.1 | 37,800 | 3.8 | 3.1-4.5 | 76,000 | 7.6 | 6.6-8.6 | 998,600 |
| 2009 | 704,700 | 70.5 | 68.8-72.1 | 177,400 | 17.7 | 16.3-19.1 | 44,200 | 4.4 | 3.6-5.2 | 73,500 | 7.4 | 6.5-8.2 | 999,800 |
| 2008 | 678,000 | 68.3 | 66.6-69.9 | 175,500 | 17.7 | 16.3-19.1 | 46,800 | 4.7 | 4.0-5.5 | 92,400 | 9.3 | 8.3-10.4 | 992,700 |
| 2007 | 679,800 | 69.3 | 67.7-71.0 | 171,900 | 17.5 | 16.2-18.8 | 43,300 | 4.4 | 3.7-5.2 | 85,300 | 8.7 | 7.7-9.7 | 980,300 |
| 2006 | 681,100 | 70.8 | 69.2-72.4 | 164,500 | 17.1 | 15.8-18.4 | 43,000 | 4.5 | 3.7-5.3 | 73,300 | 7.6 | 6.7-8.5 | 962,000 |
| 2005 | 656,500 | 67.9 | 66.3-69.5 | 187,600 | 19.4 | 18.0-20.8 | 40,500 | 4.2 | 3.6-4.8 | 81,900 | 8.5 | 7.5-9.5 | 966,600 |
| 2004 | 604,600 | 67.1 | 64.2-70.1 | 175,100 | 19.4 | 16.9-21.9 | 41,300 | 4.6 | 3.2-6.0 | 79,500 | 8.8 | 7.1-10.5 | 900,500 |
| 2003 | 737,000 | 80.4 | 78.9-81.8 | 99,600 | 10.9 | 9.7-12.0 | 27,400 | 3.0 | 2.3-3.7 | 52,900 | 5.8 | 4.9-6.6 | 917,000 |
| 2002 | 695,300 | 77.3 | 75.9-78.8 | 106,700 | 11.9 | 10.7-13.1 | 41,600 | 4.6 | 4.0-5.3 | 55,800 | 6.2 | 5.4-7.0 | 899,400 |
| 2001 | 747,500 | 84.2 | 82.6-85.7 | 80,400 | 9.1 | 7.8-10.3 | 22,600 | 2.5 | 1.8-3.3 | 37,600 | 4.2 | 3.5-5.0 | 888,200 |
| 2000 | 639,000 | 73.4 | 71.9-74.9 | 136,300 | 15.7 | 14.4-16.9 | 40,700 | 4.7 | 4.0-5.4 | 54,500 | 6.3 | 5.4-7.1 | 870,600 |

Source: Hawai'i State, Department of Health, Office of Health Status Monitoring. Hawaii Health Data Warehouse. Behavioral Risk Factor Surveillance System. Health-mental bad past 30 days