

Table 7.182 updated

Native Hawaiian Depressive Disorders in Hawai'i: 2006, 2008, 2010-2014

Survey Question: (Ever told) you have a depressive disorder, including depression, major depression, dysthymia, or minor depression?

| Year | Native Hawaiian | | | | | | Estimated Population |
|------|-----------------|------|------------|---------|------|------------|----------------------|
| | Yes | | | No | | | |
| | No. | % | C.I. (95%) | No. | % | C.I. (95%) | No. |
| 2014 | 20,900 | 15.8 | 12.5-19.2 | 111,300 | 84.2 | 80.8-87.5 | 132,200 |
| 2013 | 15,600 | 13.7 | 11.0-16.5 | 98,100 | 86.3 | 83.5-89.0 | 113,800 |
| 2012 | 14,500 | 11.5 | 8.3-14.8 | 111,500 | 88.5 | 85.2-91.7 | 126,000 |
| 2011 | 12,000 | 10.8 | 7.7-14.0 | 98,600 | 89.2 | 86.0-92.3 | 110,600 |
| 2010 | 10,600 | 8.5 | 5.5-11.6 | 113,300 | 91.5 | 88.4-94.5 | 123,900 |
| 2008 | 14,200 | 10.5 | 7.7-13.4 | 120,300 | 89.5 | 86.6-92.3 | 134,500 |
| 2006 | 9,300 | 8.3 | 5.8-10.8 | 102,500 | 91.7 | 89.2-94.2 | 111,800 |

| Year | State of Hawai'i | | | | | | Estimated Population |
|------|------------------|------|------------|---------|------|------------|----------------------|
| | Yes | | | No | | | |
| | No. | % | C.I. (95%) | No. | % | C.I. (95%) | No. |
| 2014 | 118,700 | 10.7 | 9.8-11.7 | 989,500 | 89.3 | 88.3-90.2 | 1,108,300 |
| 2013 | 125,100 | 11.4 | 10.4-12.4 | 971,000 | 88.6 | 87.6-89.6 | 1,096,100 |
| 2012 | 123,400 | 11.5 | 10.4-12.6 | 950,300 | 88.5 | 87.4-89.6 | 1,073,700 |
| 2011 | 111,000 | 10.6 | 9.5-11.7 | 936,900 | 89.4 | 88.3-90.5 | 1,047,900 |
| 2010 | 85,400 | 8.9 | 8.0-9.9 | 871,600 | 91.1 | 90.1-92.0 | 957,000 |
| 2008 | 94,500 | 9.8 | 8.9-10.8 | 866,300 | 90.2 | 89.2-91.1 | 960,700 |
| 2006 | 81,900 | 8.8 | 8.0-9.7 | 843,800 | 91.2 | 90.3-92.0 | 925,700 |

Source: Hawai'i State, Department of Health, Office of Health Status Monitoring. Hawaii Health Data Warehouse. Behavioral Risk Factor Surveillance System. Depressive disorder, for the State of Hawai'i.