

Table 7.232

Age-Adjusted Mortality Rates (AAMR) and Years of Potential Life Lost (YPLL) by Cause of Resident Deaths in Hawai'i: Aggregated Years 2002-2013

Cause of Death	State of Hawai'i Resident Deaths											
	2002-2004			2005-2007			2008-2010			2011-2013		
	Avg. Ann. # Deaths	Avg. Ann. AAMR	Avg. Ann. YPLL	Avg. Ann. # Deaths	Avg. Ann. AAMR	Avg. Ann. YPLL	Avg. Ann. # Deaths	Avg. Ann. AAMR	Avg. Ann. YPLL	Avg. Ann. # Deaths	Avg. Ann. AAMR	Avg. Ann. YPLL
Major cardiovascular diseases	3,395	82.7	16,550	3,089	68	16,070	3,073	61.8	15,610	3,157	58.6	15,980
Malignant neoplasms	2,020	49.7	16,240	2,144	49.2	17,210	2,180	46.3	16,930	2,256	44.5	17,730
All other diseases (Residual)	742	18.1	4,950	1,027	22.5	6,500	1,195	23.3	5,590	1,367	24.5	5,700
Accidents (unintentional injuries)	386	9.9	9,290	440	10.7	10,060	403	9.3	9,600	450	9.8	9,790
Chronic lower respiratory diseases	283	6.9	1,060	290	6.4	1,180	292	6	1,060	299	5.7	1,190
Influenza/Pneumonia	237	5.7	730	229	4.9	650	277	5.4	1,180	388	6.9	1,240
Diabetes mellitus	196	4.8	1,290	258	5.8	1,650	287	6	1,860	266	5.1	1,920
Alzheimer's disease	155	3.7	80	211	4.3	50	215	3.9	50	234	3.9	70
Nephritis, nephrotic syndrome and nephrosis	139	3.4	800	166	3.7	720	206	4.2	950	118	2.2	450
Symptoms, signs & abnorm clinical/lab findings NEC	145	3.6	1,540	162	3.5	1,690	128	2.6	1,330	176	3.2	1,200
Intentional self-harm (suicide)	120	3.2	3,580	118	3	3,530	167	4.1	5,100	170	3.9	5,300
Other diseases of respiratory system	137	3.3	720	138	3	580	145	3	710	154	2.9	720
Septicemia	158	3.9	1,190	159	3.6	1,160	127	2.7	1,090	119	2.3	740
Pneumonitis due to solids and liquids	118	2.8	290	129	2.7	280	115	2.1	170	104	1.8	210
Chronic liver disease and cirrhosis	79	2	1,070	94	2.2	1,470	107	2.4	1,620	124	2.5	1,720
Parkinson's disease	87	2.1	90	89	1.9	80	86	1.7	100	97	1.8	90
In situ & benign neoplasms; unknown neoplasms	54	1.3	300	55	1.2	330	72	1.5	460	70	1.3	380
Certain conditions originating in perinatal period	65	1.8	4,810	63	1.7	4,590	59	1.5	4,400	62	1.6	4,610
Events of undetermined intent	37	1	1,110	72	1.8	2,030	54	1.3	1,470	44	1	1,220
Viral hepatitis	25	0.6	440	35	0.8	580	38	0.8	620	36	0.7	600
Congenital malform/deform & chromosomal abnorms	34	0.9	1,830	31	0.8	1,640	33	0.8	1,590	31	0.7	1,560

Other & unspecified infectious & parasitic disease	30	0.7	360	32	0.7	430	28	0.6	210	21	0.4	160
Assault (homicide)	28	0.8	1,040	25	0.6	980	25	0.6	860	24	0.6	840
Infections of Kidney	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	87	1.7	420
Other disorders of circulatory system	15	0.4	150	16	0.4	130	28	0.6	140	22	0.4	230
Anemias	17	0.4	70	21	0.5	160	14	0.3	70	17	0.3	40
Peptic ulcer	15	0.4	70	15	0.3	80	15	0.3	40	16	0.3	100
Cholelithiasis and other disorders of gallbladder	9	0.2	20	10	0.2	60	14	0.3	50	18	0.4	100
Nutritional deficiencies	12	0.3	40	14	0.3	30	13	0.3	50	9	0.2	80
Complications of medical and surgical care	7	0.2	70	11	0.2	120	8	0.2	100	10	0.2	70
Certain other intestinal infections	n/r	n/r	n/r	n/r	n/r	n/r	11	0.2	90	13	0.2	70
Tuberculosis	n/r	n/r	n/r	n/r	n/r	n/r	6	0.1	20	n/r	n/r	n/r
Hernia	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r
Pregnancy, childbirth and the puerperium	n/r	n/r	n/r	6	0.2	270	n/r	n/r	n/r	n/r	n/r	n/r
Note: n/r = Not Reportable, where number of deaths between 1 and 5; AAMR = Age-Adjusted Mortality Rate per 100,000 population; YPLL = Years of Potential Life Lost to age 75 years (rounded to the nearest 10)												

Source: Hawai‘i State, Department of Health, Office of Health Status Monitoring. Hawaii Health Data Warehouse. Behavioral Risk Factor Surveillance System. Leading Causes of Death in Hawai‘i.