Table 7.52a

Five Leading Cancer Sites for Morbidity by Males and Race/Ethnicity, Hawai'i, 1995-2000.

Morbidity	Race/Ethnicity									
	Caucasian		Chinese		Filipino		Japanese		Native Hawaiian	
Male	Deaths	% Total	Deaths	% Total	Deaths	% Total	Deaths	% Total	Deaths	% Total
Colon and Rectum	148	9.1	27	7.6	77	8.9	233	12.9	93	10.8
Liver			23	6.4	55	6.4			43	5.0
Lung and Bronchus	453	27.8	122	34.2	263	30.4	441	24.5	295	34.3
NHL	74	4.5								
Pancreas	79	4.8	28	7.8	51	5.9	120	6.7		
Prostate	209	12.8	36	10.1	113	13.0	163	9.1	71	8.2
Stomach							227	12.6	57	6.6
Total, All Sites	1,630	100.0	357	100.0	866	100.0	1,800	100.0	861	100.0

Most frequently diagnosed or leading cause of cancer death

Second most frequently diagnosed or second leading cause of cancer death

Third most frequently diagnosed or third leading cause of cancer death

Note: Counts are 6 year totals. NHL = Non-Hodgkin lymphomas. Liver includes hepatic bile duct.

Source: Hawai'i State, University of Hawai'i, Cancer Research Center of Hawai'i, Hawai'i Tumor Registry, *Hawai'i Cancer Facts & Figures 2003-2004*. (Honolulu)