

Table 7.58a

Asthma by Age for Native Hawaiians in Hawai'i: 2000-2010

| Year | Native Hawaiian | | | | | | | | | |
|------|-----------------|-------|-------|-------|-------|-------|-------|-------|-------|--------|
| | Age in Years | | | | | | | | | Total |
| | 0-14 | 15-17 | 18-24 | 25-34 | 35-44 | 45-54 | 55-64 | 65-74 | 75+ | |
| 2010 | 15,402 | 3,543 | 5,993 | 5,229 | 5,478 | 4,108 | 3,502 | 1,918 | 571 | 45,744 |
| 2009 | 20,316 | 3,572 | 5,822 | 4,580 | 5,230 | 7,093 | 3,262 | 2,630 | 435 | 52,940 |
| 2008 | 16,135 | 3,858 | 6,122 | 4,576 | 5,949 | 5,373 | 4,400 | 1,852 | 974 | 49,239 |
| 2007 | 20,847 | 3,274 | 6,962 | 4,685 | 3,712 | 6,945 | 3,973 | 1,877 | 814 | 53,089 |
| 2006 | 18,754 | 3,222 | 4,186 | 5,479 | 3,584 | 4,250 | 2,742 | 1,916 | 1,161 | 45,294 |
| 2005 | 15,948 | 2,367 | 4,052 | 4,345 | 4,125 | 4,603 | 2,954 | 1,532 | 395 | 40,321 |
| 2004 | 15,165 | 2,287 | 4,329 | 4,497 | 4,560 | 2,776 | 3,320 | 1,448 | 717 | 39,099 |
| 2003 | 13,636 | 2,051 | 3,984 | 3,151 | 5,968 | 4,567 | 2,036 | 1,457 | 271 | 37,121 |
| 2002 | 13,725 | 1,749 | 3,755 | 2,666 | 3,458 | 2,056 | 1,233 | 1,747 | 1,095 | 31,484 |
| 2001 | 15,245 | 2,864 | 4,098 | 5,619 | 3,886 | 3,209 | 1,998 | 1,278 | * | 38,309 |
| 2000 | 15,575 | 2,532 | 4,067 | 2,550 | 4,284 | 2,934 | 1,607 | 1,456 | 562 | 35,567 |

Table 7.58b

Asthma by Age in Hawai'i: 2000-2010

| Year | State of Hawai'i | | | | | | | | | |
|------|------------------|-------|--------|--------|--------|--------|--------|--------|-------|---------|
| | Age in Years | | | | | | | | | Total |
| | 0-14 | 15-17 | 18-24 | 25-34 | 35-44 | 45-54 | 55-64 | 65-74 | 75+ | |
| 2010 | 34,373 | 9,127 | 16,519 | 17,516 | 15,582 | 21,395 | 17,283 | 9,427 | 6,406 | 147,628 |
| 2009 | 39,492 | 9,617 | 13,963 | 16,026 | 13,846 | 21,212 | 17,602 | 10,134 | 7,755 | 149,647 |
| 2008 | 32,873 | 8,590 | 15,088 | 13,160 | 14,853 | 22,052 | 18,893 | 7,687 | 6,916 | 140,112 |
| 2007 | 34,047 | 7,969 | 14,091 | 12,771 | 12,113 | 19,688 | 18,075 | 7,111 | 5,761 | 131,626 |
| 2006 | 40,073 | 7,435 | 12,171 | 14,475 | 14,282 | 19,092 | 14,182 | 8,386 | 6,332 | 136,428 |
| 2005 | 32,963 | 7,932 | 11,444 | 11,950 | 14,042 | 17,259 | 14,750 | 7,157 | 5,310 | 122,807 |
| 2004 | 31,786 | 6,656 | 12,280 | 14,898 | 13,696 | 14,194 | 12,468 | 6,916 | 4,246 | 117,140 |
| 2003 | 29,911 | 5,674 | 9,415 | 11,538 | 13,561 | 16,668 | 9,660 | 5,286 | 4,679 | 106,392 |
| 2002 | 26,776 | 6,328 | 12,665 | 10,685 | 12,529 | 12,409 | 9,000 | 7,594 | 4,502 | 102,488 |
| 2001 | 32,812 | 6,564 | 9,991 | 13,062 | 14,129 | 14,048 | 6,711 | 5,585 | 3,358 | 106,260 |
| 2000 | 30,798 | 7,728 | 9,006 | 9,853 | 12,789 | 13,580 | 6,372 | 5,285 | 4,580 | 99,991 |

* Number of cases in sample numerator or denominator too small for reliability.

Source: Hawai'i State, Department of Health, Hawai'i Health Survey.