

Table 7.82b

Native Hawaiian Men Aged 40 and Over in Hawai'i Who Had a Digital Rectal Exam (DRE) and Prostate Specific Antigen (PSA) Test in Hawai'i: 2005-2011

Year	Native Hawaiian Men Aged 40+							
	Percentage Who Had a Digital Rectal Exam				Percentage Who Had a Prostate Specific Antigen Test			
	Ever had a DRE		Last DRE within Past Year		Ever had a PSA Test		Last PSA Test within Last Year	
	%	C.I. (95%)	%	C.I. (95%)	%	C.I. (95%)	%	C.I. (95%)
2011	(X)		(X)		35.1	(27.0-44.1)	20.2	(13.8-28.6)
2010	58.0	(48.3-67.2)	22.7	(16.4-30.6)	49.0	(39.9-58.2)	30.2	(22.6-39.2)
2009	53.4	(40.3-66.0)	15.6	( 7.9-28.2)	37.2	(25.7-50.4)	25.6	(15.3-39.5)
2008	53.7	(44.2-62.9)	28.1	(20.1-37.7)	40.2	(31.6-49.5)	31.7	(23.3-41.5)
2007	65.2	(56.4-73.0)	28.5	(21.2-37.1)	46.1	(37.2-55.2)	27.6	(20.8-35.7)
2006	49.3	(40.4-58.2)	17.4	(12.2-24.2)	28.0	(21.3-36.0)	19.7	(14.0-27.1)
2005	66.0	(49.7-79.3)	24.8	(15.0-38.1)	32.7	(21.3-46.5)	21.7	(13.0-34.1)
(X) = Not reported								

**Source:** Hawai'i State, Department of Health, Behavioral Risk Factor Surveillance System.